

National Library PubMed of Medicine

PubMed

Nucleotide

Protein Geno

Structure

PopSet

Taxonomy

Go

ОМІМ

Search | PubMed

for

Preview/Index

History

Clipboard

Clear

About Entrez

...Display

Limits

Abstract

Save Text

Order |

Add to Clipboard

Entrez PubMed Overview Help | FAQ New/Noteworthy

PubMed Services
Journal Browser
MeSH Browser
Single Citation Matcher
Batch Citation Matcher
Clinical Queries

Related Resources Order Documents Grateful Med Consumer Health Clinical Alerts ClinicalTrials.gov

Privacy Policy

Cubby NEW

1: Arterioscler Thromb Vasc Biol 2000 Feb;20 (2):435-42

Related Articles, Books,

LinkOut

Full text article at atvb.ahajournals.org

Complete atherosclerosis regression after human ApoE gene transfer in ApoE-deficient/nude mice.

Desurmont C, Caillaud JM, Emmanuel F, Benoit P, Fruchart JC, Castro G, Branellec D, Heard JM, Duverger N

Laboratoire RTG, Institut Pasteur, Paris, France.

The apolipoprotein E (apoE)-deficient mouse is a relevant animal model of human atherosclerosis. Although the prevention of atherosclerosis development has been documented after somatic gene transfer into animal models, regression of lesions remains to be demonstrated. Thus, we used this genetically defined mouse model nn the nude background to show atherosclerosis regression. ApoE-deficient nude mice were infected with 5 x 10(8) or 10(9) plaque-forming units of a first-generation adenovirus encoding human apoE cDNA. The secretion of human apoE resulted in a rapid decrease of total cholesterol, which normalized the hypercholesterolemic phenotype within 14 days (from 600+/-100 to <100 microg/mL). Transgene expression was observed during a period of >4 months, with a normalization of cholesterol and triglyceride levels during 5 months. At that time, we successfully reinjected the recombinant adenovirus and observed the appearance of the human protein as well as the correction of lipoprotein phenotype. In mice killed 6 months-after the first infection, we observed a dose-dependent regression of fatty streak lesions in the aorta. We showed sustained expression of a transgene with a first-generation adenoviral vector and a correction of dyslipoproteinemia phenotype leading to lesion regression. These data demonstrate that somatic gene transfer can induce plaque regression.

PMID: 10669641, UI: 20136123

Display Abstract



Order Add to Clipboard

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer



National Library PubMed of Medicine

PubMed

Nucleotide

Search PubMed

Protein Gen

Structure

PopSet

Taxonomy

MIMO

for | Limits

Preview/Index

History

Clipboard

Clear

About Entrez

Entrez PubMed Overview Help | FAQ New/Noteworthy

PubMed Services
Journal Browser
MeSH Browser
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
Cubby NEW

Related Resources Order Documents Grateful Med Consumer Health Clinical Alerts ClinicalTrials.gov

Privacy Policy

Display Abstract

Save! Text

Order Add to Clipboard

1: Curr Opin Lipidol 2000 Feb;11(1):25-9

Related Articles, Books, LinkOut

The value of apolipoprotein E knockout mice for studying the effects of dietary fat and cholesterol on atherogenesis.

Osada J. Joven J., Maeda N

Departamento de Bioquimica y Biologia Molecular y Celular, Facultad de Veterinaria, Universidad de Zaragoza, Spain. Josada@posta.unizar.es

The ability of the apolipoprotein E-deficient mouse to develop spontaneous atherosclerosis, which resembles the human process, is an excellent model in which to assess the impact of dietary factors. This review discusses the role of several nutrients in the development of atherosclerosis and the mechanisms through which they act.

Publication Types:

- Review
- Review, tutorial

PMID: 10750690, UI: 20212728

Display Abstract

Save: Text

Order Add to Clipboard

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer